*(*'

arylsulfinyl, arylsulfonyl, thio, alkylthio, alkylsulfinyl, alkylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, halogen, haloalkyl, polyhaloalkyl, polyhaloalkoxy, aminothio, aminosulfinyl, aminosulfonyl, alkyisulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, heteroarylsulfonylamino, acyloarbonylamino, arylcarbonyloxy, arylcarbonylamino, heteroarylcarbonyloxy, heteroarylcarbonylamino, cyano, nitro, alkenylcarbonylamino, alkynylcarbonylamino, alkylaminocarbonylamino, arylaminocarbonylamino, alkenylaminocarbonylamino, alkynyloxycarbonylamino, alkoxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, aryloxycarbonylamino, heteroaryloxycarbonylamino, alkylaminocarbonyloxy, alkoxycarbonylamino, l,l-(alkoxyl or aryloxy)<sub>2</sub>alkyl (where the two aryl or alkyl substituents can be independently defined, or linked to one another to form a ring), S(O)<sub>2</sub>R<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>(C=NR<sup>7</sup>)alkyl, -NR<sup>6</sup>(C=NR<sup>7</sup>)alkenyl,

-NR<sup>6</sup>(C=NR<sup>7</sup>)alkynyl, -NR<sup>6</sup>(C=NR<sup>7</sup>)heteroaryl, -NR<sup>8</sup>(C=NCN)-amino,

pyridine-N-oxide,

$$-N \longrightarrow R^8 \longrightarrow N \longrightarrow O \longrightarrow N$$

(where Q is O or H<sub>2</sub> and n' is 0, 1, 2 or 3) or

—C=CH—C—R<sup>8a</sup>; -PO(R<sup>13</sup>)(R<sup>14</sup>), (where R<sup>13</sup> and R<sup>14</sup> are independently alkyl, aryl, alkoxy, aryloxy, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy, cycloheteroalkyl, cycloheteroalkoxy, or cycloheteroalkylalkoxy);

R6, R7, R8 and R9 are the same or different and are independently hydrogen, alkyl, haloalkyl, aryl, heteroaryl, arylalkyl, cycloalkyl, (cycloalkyl)alkyl, or cycloheteroalkyl;

and R<sup>1</sup> may be unsubstituted or substituted with from one to five substituents; and wherein the R<sup>1</sup> heteroaryl group is selected from

R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are the same or different and are independently H, alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, (alkyl or aryl) Si (where each alkyl or aryl group is independent), cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, alkenylamino, alkynylamino, arylalkylamino, aryl, arylalkyl, arylamino, aryloxy, cycloheteroalkyl, cycloheteroalkylalkyl, heteroaryl, heteroarylamino, heteroaryloxy, arylthio, arylsulfinyl, arylsulfonyl, thio, alkylthio, alkylsulfinyl, alkylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, halogen, haloalkyl, polyhaloalkoxy, aminothio, aminosulfinyl, aminosulfonyl, alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, hydroxy, acyl, carboxy, aminocarbonyl, alkylcarbonyl, alkylcarbonylamino, arylcarbonyl, arylcarbonyloxy, arylcarbonylamino, heteroarylcarbonyl, heteroarylcarbonylamino, arylcarbonylamino, cyano, nitro, alkenylcarbonylamino, alkynylaminocarbonylamino, arylaminocarbonylamino, alkynylaminocarbonylamino, arylaminocarbonylamino, alkynylaminocarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, alkynyloxycarbonyla

an inocarbonylamino, alkylaminocarbonyloxy, alkoxycarbonylamino, I,I-(alkoxyl or aryloxy)2alkyl (where the two aryl or alkyl substituents can be independently defined, or linked to one another to form a ring),  $S(O)_2R^6R^7$ ,  $-NR^6(C=NR^7)_2kyl$ ,  $-NR^6(C=NR^7)_2kyl$ ,  $-NR^6(C=NR^7)_2kyl$ ,

-NR $^6$ (C=N $^7$ )alkynyl, -NR $^6$ (C=NR $^7$ )heteroaryl, -NR $^8$ (C=NCN)-amino,

pyridine-N-oxide,

$$-N \longrightarrow R^{8} \longrightarrow N \longrightarrow N \longrightarrow N$$


(where Q is O or H₂ and n' is 0, 1,2 or 3) or


—C=CH—C—R<sup>8a</sup>; tetrazolyl, pyrazolyl, pyridyl, thiazolyl, pyrimidinyl, imidazole, oxazole, or triazole, -PO(R<sup>13</sup>)(R<sup>14</sup>), (where R<sup>13</sup> and R<sup>14</sup> are independently alkyl, aryl, alkoxy, aryloxy, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, cycloheteroalkylalkoxy, and may be optionally independently substituted with from one to five substituents, which may be the same or different;

including pharmaceutically acceptable salts thereof, prodrugs thereof, and all stereoisomers thereof. --

--14. (Amended) The compound as defined in Claim I wherein the moiety


75.b







--19. (Twice Amended) The compound as defined in Claim I wherein the R¹ heteroaryI

group is










--22. (Twice Amended) The compound as defined in Claim 14 wherein

$$R^{1}$$
 is  $R_{1}$   $R_{2}$   $R_{3}$   $R_{4}$   $R$ 

--24. (Twice Amended) The compound as defined in Claim 14 wherein

28 50 h



--64. (Amended) A compound having the structure

5 Jy

wherein n is 4;

X is N;

Z is a heteroaryl group;

R¹ is tetrazolyl, pyrazolyl, thiazolyl, pyrimidinyl, , oxazole, or triazole

R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are the same or different and are independently hydrogen, alkyl, haloalkyl, aryl, heteroaryl, arylalkyl, cycloalkyl, (cycloalkyl)alkyl, or cycloheteroalkyl;

and R<sup>1</sup> may be unsubstituted or substituted with from one to five substituents;

R<sup>2</sup>\ R<sup>3</sup> and R<sup>4</sup> are the same or different and are independently H, alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, alkynyloxy, (alkyl or aryl)3Si (where each alkyl or aryl group is independent), cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, alkenylamino, alkynylamino, arylalkylamino, aryl, arylalkyl, arylamino, aryloxy, cycloheteroalkyl, cycloheteroalkylalkyl, heteroaryl, heteroarylamino,\heteroaryloxy, arylthio, arylsulfinyl, arylsulfonyl, thio, alkylthio, alkylsulfinyl, alkylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, halogen, haloalkyl, polyhaloalkyl, polyhaloalkoxy, aminothio, aminosulfinyl, aminosulfonyl, alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, hydroxy, acyl, carboxy, aminocarbonyl, alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylcarbonylamino, arylcarbonyl, arylcarbonyloxy, arylcarbonylamino, heteroarylcarbonyl, heteroarylcarbonyloxy, heteroarylcarbonylamino, cyano, nitro, alkenylcarbonylamino, àlkynylcarbonylamino, alkylaminocarbonylamino, alkenylaminocarbonylamino, alkynylaminocarbonylamino, arylaminocarbonylamino, heteroarylaminocarbonylamino, àlkoxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, aryloxycarbonylamino, heteroaryloxycarbonylamino, aminocarbonylamino, alkylaminocarbonyloxy, alkoxycarbonylamino, I,I-(alkoxyl or aryloxy)2alkyl (where the two aryl or alkyl substituents can be independently defined, or linked to one another to form a ring),  $S(O)_2R^6R^7$ ,  $-NR^6(C=NR^7)_alkyl$ ,  $-NR^6(C=NR^7)_alkenyl$ ,

-NR<sup>6</sup>(C=NR<sup>7</sup>)alkynyl, -NR<sup>6</sup>(C=NR<sup>7</sup>)heteroaryl, -NR<sup>8</sup>(C=NCN)-amino,

$$-\frac{\overset{\text{O}}{\underset{\text{P}}{\bigvee}}\overset{\text{O}}{\underset{\text{O}}{\longleftarrow}} \mathbb{R}^8}{\overset{\text{R}}{\underset{\text{O}}{\longleftarrow}}}$$

pyridine-N-oxide,

$$-N \longrightarrow R^8 \longrightarrow N \longrightarrow N \longrightarrow N$$

(where Q is O or  $H_2$  and n' is 0, 1, 2 or 3) or

NR<sup>8</sup>R<sup>9</sup> O | C=CH-C-R<sup>8a</sup>; tetrazolyl, pyrazolyl, pyridyl, thiazolyl, pyrimidinyl, imidazole, oxazole, or triazole, -PO(R<sup>13</sup>)(R<sup>14</sup>), (where R<sup>13</sup> and R<sup>14</sup> are independently alkyl, aryl, alkoxy, aryloxy, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl,